



# St. Vincent Healthcare

1233 North 30<sup>th</sup> Street, Billings, Montana 59101

## STROKE-INFARCTION/RULE OUT STROKE-INFARCTION TRANSIENT ISCHEMIC ATTACK ADMISSION ORDERS

1. Inpatient admission to 4T monitored 3T monitored ICU for Dr. \_\_\_\_\_
2. May go to procedures unmonitored
3. Anti-anxiety on call to MRI: Ativan 0.5 to 1 mg IV/IM( if NPO) or PO  
Valium 5 mg IV/IM (if NPO) or PO  
Other \_\_\_\_\_
4. Pulmonologist/Intensivist Consult
5. Notify Stroke Coordinator at 7668
6. Primary diagnosis Stroke-Infarction: ORDER PT and OT CONSULT TIA Rule Out Stroke
7. Code Status No CPR Full Code Other \_\_\_\_\_
8. Allergies \_\_\_\_\_
9. Height and Weight on admission to floor.
10. 0.9 NS @ 70 cc/hour \_ TKO \_ Saline lock when taking po
11. Vital signs/neuro checks per protocol Vital signs/neuro checks other \_\_\_\_\_  
Initiate policies/protocols for management of blood pressure, temperature, glucose, aspiration precautions, fall precautions, etc.
12. NPO DIET type: \_\_\_\_\_  
NPO/SLP (Speech/Language) Consult –may take medications in pureed consistency  
NPO/SLP (Speech/Language) Consult –hold medications
13. Fasting lipid panel CBC, platelets, BMP, PT/INR, PTT if not done in ED CPK with Troponins on admit  
HbA1C Antiphospholipid Syndrome Evaluator Protein C and S Antithrombin III screening  
Homocysteine Factor-V Leiden BMP daily x2 Hct/Hgb daily x2
14. Carotid Duplex US MRI Brain MRA Brain (Circle of Willis) MRA neck CT head @ 24 hours post-thrombolysis  
CTA Brain (Circle of Willis) CTA neck CT head @ 24 hours post-thrombolysis  
Echocardiogram (TTE) - suggested for patients with known cardiac disease to evaluate LVEF, valves, dyskinesias  
TEE – Patient must be NPO. Suggested for patients without known cardiac sources of emboli or negative TTE
15. BGs ac and hs x36 hours; BGs q8 hours x36 hours if NPO  
Sliding scale insulin: For BG 150-200mg/dl, give 4 units regular insulin subc  
For BG 201-250mg/dl, give 8 units regular insulin subc  
For BG 251-300mg/dl, give 12 units regular insulin subc  
For BG >300mg/dl, give 18 units regular insulin subc  
If BG >300mg/dl x2 notify Dr. \_\_\_\_\_
16. Bedrest with head of bed 30-45° Bedrest with bathroom privileges Up with assist Up ad lib
17. Thigh high support hose Pneumatic thigh highs Lovenox 40mg subc daily start \_\_\_\_\_
18. Respiratory Care Consult Rehabilitation Medicine consult
19. EC aspirin \_\_\_\_\_mg po daily Plavix \_\_\_\_\_mg po daily  
Aggrenox 1 cap po bid Pepcid 20mg IVq 12 hours; oral route when taking po Colace 1 cap po bid
20. Other Medications  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_